



PLI/GB 2003 / 0 0 4 7 6 6  
29 APR 2005



INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road

Newport  
South Wales

NP10 8QQ

REC'D 11 DEC 2003

WPO

PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

*Stephen Hordley*

Dated

17 November 2003

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)



Patents Act 1977  
(Rule 16)

13MAY03 E806777-1 002934  
P01/7700 0.00-0310932.9

The Patent Office

Cardiff Road  
Newport  
South Wales  
NP10 8QQ

# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

1. Your reference 100891-2

2. Patent application number  
(The Patent Office will fill in this part)

13 MAY 2003

0310932.9

3. Full name, address and postcode of the or of each applicant (underline all surnames)

AstraZeneca AB  
SE-151 85 Sodertalje  
Sweden

Patents ADP number (if you know it) 07822448003

If the applicant is a corporate body, give the country/state of its incorporation

Sweden

4. Title of the invention

CHEMICAL COMPOUNDS

5. Name of your agent (if you have one)

Lucy PADGET

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

AstraZeneca UK Limited  
Global Intellectual Property  
Mereside, Alderley Park  
Macclesfield,  
Cheshire SK10 4TG

Patents ADP number (if you know it)

08501132001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number  
(if you know it)

Date of filing  
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
  - b) there is an inventor who is not named as an applicant, or
  - c) any named applicant is a corporate body.
- See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description

43

Claim(s)

2

Abstract

1

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents  
(*please specify*)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Jennifer Bennett, Authorised Signatory

Date 12 May 2003

12. Name and daytime telephone number of person to contact in the United Kingdom

Shirley Douglas - 01625 510057

**Warning**

*After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.*

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.*
- Write your answers in capital letters using black ink or you may type them.*
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.*
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.*
- Once you have filled in the form you must remember to sign and date it.*
- For details of the fee and ways to pay please contact the Patent Office.*

CHEMICAL COMPOUNDS

This invention relates to chemical compounds, or pharmaceutically acceptable salts thereof. These compounds possess human 11- $\beta$ -hydroxysteroid dehydrogenase type 1 enzyme (11 $\beta$ HSD1) inhibitory activity and accordingly have value in the treatment of disease states including metabolic syndrome and are useful in methods of treatment of a warm-blooded animal, such as man. The invention also relates to processes for the manufacture of said compounds, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments to inhibit 11 $\beta$ HSD1 in a warm-blooded animal, such as man.

10        Glucocorticoids (cortisol in man, corticosterone in rodents) are counter regulatory hormones i.e. they oppose the actions of insulin (Dallman MF, Strack AM, Akana SF et al. 1993; Front Neuroendocrinol 14, 303-347). They regulate the expression of hepatic enzymes involved in gluconeogenesis and increase substrate supply by releasing glycerol from adipose tissue (increased lipolysis) and amino acids from muscle (decreased protein synthesis and  
15 increased protein degradation). Glucocorticoids are also important in the differentiation of pre-adipocytes into mature adipocytes which are able to store triglycerides (Bujalska IJ et al. 1999; Endocrinology 140, 3188-3196). This may be critical in disease states where glucocorticoids induced by "stress" are associated with central obesity which itself is a strong risk factor for type 2 diabetes, hypertension and cardiovascular disease (Bjorntorp P &  
20 Rosmond R 2000; Int. J. Obesity 24, S80-S85)

It is now well established that glucocorticoid activity is controlled not simply by secretion of cortisol but also at the tissue level by intracellular interconversion of active cortisol and inactive cortisone by the 11-beta hydroxysteroid dehydrogenases, 11 $\beta$ HSD1 (which activates cortisone) and 11 $\beta$ HSD2 (which inactivates cortisol) (Sandeep TC & Walker  
25 BR 2001 Trends in Endocrinol & Metab. 12, 446-453). That this mechanism may be important in man was initially shown using carbenoxolone (an anti-ulcer drug which inhibits both 11 $\beta$ HSD1 and 2) treatment which (Walker BR et al. 1995; J. Clin. Endocrinol. Metab. 80, 3155-3159) leads to increased insulin sensitivity indicating that 11 $\beta$ HSD1 may well be regulating the effects of insulin by decreasing tissue levels of active glucocorticoids (Walker  
30 BR et al. 1995; J. Clin. Endocrinol. Metab. 80, 3155-3159).

Clinically, Cushing's syndrome is associated with cortisol excess which in turn is associated with glucose intolerance, central obesity (caused by stimulation of pre-adipocyte differentiation in this depot), dyslipidaemia and hypertension. Cushing's syndrome shows a

---

number of clear parallels with metabolic syndrome. Even though the metabolic syndrome is not generally associated with excess circulating cortisol levels (Jessop DS et al. 2001; J. Clin. Endocrinol. Metab. 86, 4109-4114) abnormally high  $11\beta$ HSD1 activity within tissues would be expected to have the same effect. In obese men it was shown that despite having similar or  
5 lower plasma cortisol levels than lean controls,  $11\beta$ HSD1 activity in subcutaneous fat was greatly enhanced (Rask E et al. 2001; J. Clin. Endocrinol. Metab. 1418-1421). Furthermore, the central fat, associated with the metabolic syndrome expresses much higher levels of  $11\beta$ HSD1 activity than subcutaneous fat (Bujalska IJ et al. 1997; Lancet 349, 1210-1213). Thus there appears to be a link between glucocorticoids,  $11\beta$ HSD1 and the metabolic  
10 syndrome.

$11\beta$ HSD1 knock-out mice show attenuated glucocorticoid-induced activation of gluconeogenic enzymes in response to fasting and lower plasma glucose levels in response to stress or obesity (Kotelevtsev Y et al. 1997; Proc. Natl. Acad. Sci USA 94, 14924-14929) indicating the utility of inhibition of  $11\beta$ HSD1 in lowering of plasma glucose and hepatic  
15 glucose output in type 2 diabetes. Furthermore, these mice express an anti-atherogenic lipoprotein profile, having low triglycerides, increased HDL cholesterol and increased apolipoprotein AI levels. (Morton NM et al. 2001; J. Biol. Chem. 276, 41293-41300). This phenotype is due to an increased hepatic expression of enzymes of fat catabolism and PPAR $\alpha$ . Again this indicates the utility of  $11\beta$ HSD1 inhibition in treatment of the  
20 dyslipidaemia of the metabolic syndrome.

The most convincing demonstration of a link between the metabolic syndrome and  $11\beta$ HSD1 comes from recent studies of transgenic mice over-expressing  $11\beta$ HSD1 (Masuzaki H et al. 2001; Science 294, 2166-2170). When expressed under the control of an adipose specific promoter,  $11\beta$ HSD1 transgenic mice have high adipose levels of  
25 corticosterone, central obesity, insulin resistant diabetes, hyperlipidaemia and hyperphagia. Most importantly, the increased levels of  $11\beta$ HSD1 activity in the fat of these mice are similar to those seen in obese subjects. Hepatic  $11\beta$ HSD1 activity and plasma corticosterone levels were normal, however, hepatic portal vein levels of corticosterone were increased 3 fold and it is thought that this is the cause of the metabolic effects in liver.

30 Overall it is now clear that the complete metabolic syndrome can be mimicked in mice simply by overexpressing  $11\beta$ HSD1 in fat alone at levels similar to those in obese man.

11 $\beta$ HSD1 tissue distribution is widespread and overlapping with that of the glucocorticoid receptor. Thus, 11 $\beta$ HSD1 inhibition could potentially oppose the effects of glucocorticoids in a number of physiological/pathological roles. 11 $\beta$ HSD1 is present in human skeletal muscle and glucocorticoid opposition to the anabolic effects of insulin on protein turnover and glucose metabolism are well documented (Whorwood CB et al. 2001; J. Clin. Endocrinol. Metab. 86, 2296-2308). Skeletal muscle must therefore be an important target for 11 $\beta$ HSD1 based therapy.

Glucocorticoids also decrease insulin secretion and this could exacerbate the effects of glucocorticoid induced insulin resistance. Pancreatic islets express 11 $\beta$ HSD1 and carbenoxolone can inhibit the effects of 11-dehydrocorticosterone on insulin release (Davani B et al. 2000; J. Biol. Chem. 275, 34841-34844). Thus in treatment of diabetes 11 $\beta$ HSD1 inhibitors may not only act at the tissue level on insulin resistance but also increase insulin secretion itself.

Skeletal development and bone function is also regulated by glucocorticoid action. 11 $\beta$ HSD1 is present in human bone osteoclasts and osteoblasts and treatment of healthy volunteers with carbenoxolone showed a decrease in bone resorption markers with no change in bone formation markers (Cooper MS et al 2000; Bone 27, 375-381). Inhibition of 11 $\beta$ HSD1 activity in bone could be used as a protective mechanism in treatment of osteoporosis.

Glucocorticoids may also be involved in diseases of the eye such as glaucoma. 11 $\beta$ HSD1 has been shown to affect intraocular pressure in man and inhibition of 11 $\beta$ HSD1 may be expected to alleviate the increased intraocular pressure associated with glaucoma (Rauz S et al. 2001; Investigative Ophthalmology & Visual Science 42, 2037-2042).

There appears to be a convincing link between 11 $\beta$ HSD1 and the metabolic syndrome both in rodents and in humans. Evidence suggests that a drug which specifically inhibits 11 $\beta$ HSD1 in type 2 obese diabetic patients will lower blood glucose by reducing hepatic gluconeogenesis, reduce central obesity, improve the atherogenic lipoprotein phenotype, lower blood pressure and reduce insulin resistance. Insulin effects in muscle will be enhanced and insulin secretion from the beta cells of the islet may also be increased.

Currently there are two main recognised definitions of metabolic syndrome.

1) The Adult Treatment Panel (ATP III 2001 JMA) definition of metabolic syndrome indicates that it is present if the patient has three or more of the following symptoms:


- Waist measuring at least 40 inches (102 cm) for men, 35 inches (88 cm) for women;
  - Serum triglyceride levels of at least 150 mg/dl (1.69 mmol/l);
  - HDL cholesterol levels of less than 40 mg/dl (1.04 mmol/l) in men, less than 50 mg/dl (1.29 mmol/l) in women;
- 5    ➤ Blood pressure of at least 135/80 mm Hg; and / or
- Blood sugar (serum glucose) of at least 110 mg/dl (6.1 mmol/l).

2) The WHO consultation has recommended the following definition which does not imply causal relationships and is suggested as a working definition to be improved upon in due course:

- 10    ➤ The patient has at least one of the following conditions: glucose intolerance, impaired glucose tolerance (IGT) or diabetes mellitus and/or insulin resistance; together with two or more of the following:
- Raised Arterial Pressure;
  - Raised plasma triglycerides
- 15    ➤ Central Obesity
- Microalbuminuria

We have found that the compounds defined in the present invention, or a pharmaceutically acceptable salt thereof, are effective 11 $\beta$ HSD1 inhibitors, and accordingly have value in the treatment of disease states associated with metabolic syndrome.

20    Accordingly there is provided the use of a compound of formula (I):



wherein:

Ring A is selected from carbocyclyl or heterocyclyl;

- 25     $R^1$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ carbamoyl,  $C_{1-4}$ alkylS(O) $_a$  wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ sulphamoyl,  $C_{1-4}$ alkylsulphonylamino,
- 30    tri-( $C_{1-4}$ alkyl)silyloxy, carbocyclyl, heterocyclyl, carbocyclyl $C_{0-4}$ alkylene-Y- and

heterocyclylC<sub>0-4</sub>alkylene-Y-; wherein R<sup>1</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>6</sup>; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>7</sup>;

n is 0-3; wherein the values of R<sup>1</sup> may be the same or different;

5 R<sup>2</sup> and R<sup>3</sup> are independently selected from hydrogen, hydroxy, amino, cyano, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, *N*-(C<sub>1-4</sub>alkyl)amino, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, carbocyclyl, heterocyclyl, carbocyclylC<sub>1-4</sub>alkyl, heterocyclylC<sub>1-4</sub>alkyl; wherein R<sup>2</sup> and R<sup>3</sup> may be independently optionally substituted on carbon by one or more groups selected from R<sup>8</sup>; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group  
10 selected from R<sup>9</sup>;

one of R<sup>4</sup> and R<sup>5</sup> is selected from C<sub>1-4</sub>alkyl and the other is selected from hydrogen or C<sub>1-4</sub>alkyl; wherein R<sup>4</sup> and R<sup>5</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>10</sup>;

Y is -S(O)<sub>a</sub>-, -O-, -NR<sup>12</sup>-, -C(O)-, -C(O)NR<sup>13</sup>-, -NR<sup>14</sup>C(O)- or -SO<sub>2</sub>NR<sup>15</sup>-; wherein a is  
15 0 to 2;

R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> are independently selected from hydrogen, phenyl and C<sub>1-4</sub>alkyl;

R<sup>6</sup> and R<sup>8</sup> are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, *N*-(C<sub>1-4</sub>alkyl)amino, 20 *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, *N*-(C<sub>1-4</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, *N*-(C<sub>1-4</sub>alkyl)sulphamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino, carbocyclyl and heterocyclyl; wherein R<sup>6</sup> and R<sup>8</sup> may be independently optionally substituted on carbon by one or more R<sup>11</sup>;

25 R<sup>10</sup> is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, *N*-(C<sub>1-4</sub>alkyl)amino, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, *N*-(C<sub>1-4</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, *N*-(C<sub>1-4</sub>alkyl)sulphamoyl, 30 *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino; wherein R<sup>10</sup> may be independently optionally substituted on carbon by one or more R<sup>16</sup>;



$R^7$  and  $R^9$  are independently selected from  $C_{1-4}$ alkyl,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkylsulphonyl,  $C_{1-4}$ alkoxycarbonyl, carbamoyl,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl;

$R^{11}$  and  $R^{16}$  are independently selected from halo, nitro, cyano, hydroxy,


- 5 trifluoromethoxy, trifluoromethyl, amino, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxymethyl, dimethylamino, diethylamino,  $N$ -methyl- $N$ -ethylamino, acetylaminomethyl,  $N$ -methylcarbamoyl,  $N$ -ethylcarbamoyl,  $N,N$ -dimethylcarbamoyl,  $N,N$ -diethylcarbamoyl,  $N$ -methyl- $N$ -ethylcarbamoyl, methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl, ethylsulphonyl, methoxycarbonyl, ethoxycarbonyl,  $N$ -methylsulphamoyl,  $N$ -ethylsulphamoyl,  $N,N$ -dimethylsulphamoyl,  $N,N$ -diethylsulphamoyl or  $N$ -methyl- $N$ -ethylsulphamoyl;

or a pharmaceutically acceptable salt thereof;

in the manufacture of a medicament for use in the inhibition of 11 $\beta$ HSD1.

According to a further feature of the invention there is provided a compound of

15 formula (Ia):



(Ia)

wherein:

Ring A is selected from phenyl, pyridyl, thiazolyl, thienyl and furyl;

- 20  $R^1$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ sulphamoyl,  $C_{1-4}$ alkylsulphonylamino; wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^6$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^7$ ;

n is 0-3; wherein the values of  $R^1$  may be the same or different;

$R^2$  and  $R^3$  are independently selected from hydrogen, hydroxy, amino, cyano,

- 30  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ amino, carbocyclyl, heterocyclyl,

carbocyclylC<sub>1-4</sub>alkyl, heterocyclylC<sub>1-4</sub>alkyl; wherein R<sup>2</sup> and R<sup>3</sup> may be independently optionally substituted on carbon by one or more groups selected from R<sup>8</sup>; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>9</sup>;

5 R<sup>4</sup> and R<sup>5</sup> are independently selected from C<sub>1-4</sub>alkyl; wherein R<sup>4</sup> and R<sup>5</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>10</sup>;

R<sup>6</sup> and R<sup>8</sup> are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, *N*-(C<sub>1-4</sub>alkyl)amino, 10 *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, *N*-(C<sub>1-4</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, *N*-(C<sub>1-4</sub>alkyl)sulphamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino; wherein R<sup>6</sup> and R<sup>8</sup> may be independently optionally substituted on carbon by one or more R<sup>11</sup>;

R<sup>10</sup> is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, 15 mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, *N*-(C<sub>1-4</sub>alkyl)amino, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, *N*-(C<sub>1-4</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, *N*-(C<sub>1-4</sub>alkyl)sulphamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino; wherein R<sup>10</sup> may be independently 20 optionally substituted on carbon by one or more R<sup>16</sup>;

R<sup>7</sup> and R<sup>9</sup> are independently selected from C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkylsulphonyl, C<sub>1-4</sub>alkoxycarbonyl, carbamoyl, *N*-(C<sub>1-4</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl;

R<sup>11</sup> and R<sup>16</sup> are independently selected from halo, nitro, cyano, hydroxy, 25 trifluoromethoxy, trifluoromethyl, amino, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxyl, methylamino, ethylamino, dimethylamino, diethylamino, *N*-methyl-*N*-ethylamino, acetylamino, *N*-methylcarbamoyl, *N*-ethylcarbamoyl, *N,N*-dimethylcarbamoyl, *N,N*-diethylcarbamoyl, *N*-methyl-*N*-ethylcarbamoyl, methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl, ethylsulphonyl, methoxycarbonyl, 30 ethoxycarbonyl, *N*-methylsulphamoyl, *N*-ethylsulphamoyl, *N,N*-dimethylsulphamoyl, *N,N*-diethylsulphamoyl or *N*-methyl-*N*-ethylsulphamoyl;

or a pharmaceutically acceptable salt thereof; with the proviso that said compound is not (*N*-methyl-*N*-butylsulphamoylmethyl)(phenyl)ketone; [1-(*N,N*-

dimethylsulphamoyl)ethyl](phenyl)ketone; (*N,N*-dimethylsulphamoylmethyl)(4-nitrophenyl)ketone; (*N,N*-dimethylsulphamoylmethyl)(4-fluoro-2-methylaminophenyl)ketone; (*N,N*-dimethylsulphamoylmethyl)(3-methoxy-4-methyl-6-aminophenyl)ketone; (*N,N*-dimethylsulphamoylmethyl)(3-methoxy-6-aminophenyl)ketone; (*N,N*-  
 5 dimethylsulphamoylmethyl)(phenyl)ketone; (*N,N*-dimethylsulphamoylmethyl)(2-nitro-4-methoxyphenyl)ketone; (*N,N*-dimethylsulphamoylmethyl)(2-amino-4-methoxyphenyl)ketone; [1-(*N*-methyl-*N*-butylsulphamoyl)ethyl](phenyl)ketone; or (*N,N*-dimethylsulphamoylmethyl)(thien-2-yl)ketone.

In this specification the term "alkyl" includes both straight and branched chain alkyl  
 10 groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. For example, "C<sub>1-4</sub>alkyl" includes propyl, isopropyl and *t*-butyl. However, references to individual alkyl groups such as 'propyl' are specific for the straight chained version only and references to individual branched chain alkyl groups such as 'isopropyl' are specific for the branched chain version only. A similar convention applies to other radicals  
 15 therefore "carbocyclylC<sub>1-4</sub>alkyl" includes 1-carbocyclylpropyl, 2-carbocyclylethyl and 3-carbocyclylbutyl. The term "halo" refers to fluoro, chloro, bromo and iodo.

Where optional substituents are chosen from "one or more" groups it is to be understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups.

20 "Heteroaryl" is a totally unsaturated, mono or bicyclic ring containing 3-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked. Suitably "heteroaryl" refers to a totally unsaturated, monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 8 - 10 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may,  
 25 unless otherwise specified, be carbon or nitrogen linked. Examples and suitable values of the term "heteroaryl" are thienyl, furyl, thiazolyl, pyrazolyl, isoxazolyl, imidazolyl, pyrrolyl, thiadiazolyl, isothiazolyl, triazolyl, pyranyl, indolyl, pyrimidyl, pyrazinyl, pyridazinyl, benzothienyl, pyridyl and quinolyl. Particularly "heteroaryl" refers to thienyl, furyl, thiazolyl, pyridyl, benzothienyl, imidazolyl or pyrazolyl.

30 "Aryl" is a totally unsaturated, mono or bicyclic carbon ring that contains 3-12 atoms. Suitably "aryl" is a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. Suitable values for "aryl" include phenyl or naphthyl. Particularly "aryl" is phenyl.

A "heterocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 3-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH<sub>2</sub>- group can optionally be replaced by a -C(O)- or a ring sulphur atom may be optionally oxidised to form the S-oxides. Preferably a "heterocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 5 or 6 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH<sub>2</sub>- group can optionally be replaced by a -C(O)- or a ring sulphur atom may be optionally oxidised to form S-oxide(s). Examples and suitable values of the term "heterocyclyl" are thienyl, piperidiny, morpholiny, furyl, thiazolyl, pyridyl, imidazolyl, 1,2,4-triazolyl, thiomorpholiny, coumariny, pyrimidinyl, phthalidyl, pyrazolyl, pyraziny, pyridaziny, benzothienyl, benzimidazolyl, tetrahydrofuryl, [1,2,4]triazolo[4,3-a]pyrimidinyl, piperidinyl, indolyl, 1,3-benzodioxolyl and pyrrolidinyl.

A "carbocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic carbon ring that contains 3-12 atoms; wherein a -CH<sub>2</sub>- group can optionally be replaced by a -C(O)-. Preferably "carbocyclyl" is a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. Suitable values for "carbocyclyl" include cyclopropyl, cyclobutyl, 1-oxocyclopentyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, phenyl, naphthyl, tetralinyl, indanyl or 1-oxoindanyl. Particularly "carbocyclyl" is cyclohexyl, phenyl, naphthyl or 2-6-dioxocyclohexyl.

An example of "C<sub>1-4</sub>alkanoyloxy" is acetoxy. Examples of "C<sub>1-4</sub>alkoxycarbonyl" include methoxycarbonyl, ethoxycarbonyl, *n*- and *t*-butoxycarbonyl. Examples of "C<sub>1-4</sub>alkoxy" include methoxy, ethoxy and propoxy. Examples of "oxyC<sub>1-4</sub>alkoxy" include oxymethoxy, oxyethoxy and oxypropoxy. Examples of "C<sub>1-4</sub>alkanoylamino" include formamido, acetamido and propionylamino. Examples of and "C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2" include methylthio, ethylthio, methylsulphiny, ethylsulphiny, mesyl and ethylsulphonyl. Examples of and "C<sub>1-4</sub>alkylsulphonyl" include mesyl and ethylsulphonyl. Examples of "C<sub>1-4</sub>alkanoyl" include propionyl and acetyl. Examples of "*N*-(C<sub>1-4</sub>alkyl)amino" include methylamino and ethylamino. Examples of "*N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino" include di-*N*-methylamino, di-(*N*-ethyl)amino and *N*-ethyl-*N*-methylamino. Examples of "C<sub>2-4</sub>alkenyl" are vinyl, allyl and 1-propenyl. Examples of "C<sub>2-4</sub>alkynyl" are ethynyl, 1-propynyl and 2-propynyl. Examples of "*N*-(C<sub>1-4</sub>alkyl)sulphamoyl" are *N*-(C<sub>1-3</sub>alkyl)sulphamoyl, *N*-(methyl)sulphamoyl and *N*-(ethyl)sulphamoyl. Examples of

"*N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl" are *N,N*-(dimethyl)sulphamoyl and *N*-(methyl)-*N*-(ethyl)sulphamoyl. Examples of "*N*-(C<sub>1-4</sub>alkyl)carbamoyl" are methylaminocarbonyl and ethylaminocarbonyl. Examples of "*N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl" are dimethylaminocarbonyl and methylethylaminocarbonyl. Examples of

5 "*C*<sub>1-4</sub>alkylsulphonylamino" are mesylamino and ethylsulphonylamino. Examples of "*C*<sub>0-4</sub>alkylene" are a direct bond, methylene and ethylene. Examples of "tri-(C<sub>1-4</sub>alkyl)silyloxy" include tri-(methyl)silyloxy dimethyl-*t*-butylsilyloxy.

A suitable pharmaceutically acceptable salt of a compound of the invention is, for example, an acid-addition salt of a compound of the invention which is sufficiently basic, for

10 example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. In addition a suitable pharmaceutically acceptable salt of a compound of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an

15 organic base which affords a physiologically-acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

Some compounds of the formula (I) may have chiral centres and/or geometric isomeric centres (*E*- and *Z*- isomers), and it is to be understood that the invention

20 encompasses all such optical, diastereoisomers and geometric isomers that possess 11 $\beta$ HSD1 inhibitory activity.

The invention relates to any and all tautomeric forms of the compounds of the formula (I) that possess 11 $\beta$ HSD1 inhibitory activity.

It is also to be understood that certain compounds of the formula (I) can exist in

25 solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess 11 $\beta$ HSD1 inhibitory activity.

Particular values of variable groups are as follows. Such values may be used where appropriate with any of the definitions, claims or embodiments defined hereinbefore or

30 hereinafter.

Ring A is selected from aryl.

Ring A is heteroaryl.

Ring A is carbocyclyl.

Ring A is heterocyclyl.

Ring A is pyridyl, phenyl, thienyl, furyl or pyrazinyl.

Ring A is pyrid-2-yl, phenyl, thien-2-yl, fur-2-yl, pyrazin-2-yl.

Ring A is pyridyl, phenyl, thienyl, furyl, pyrazinyl, 1,2,3-thiadiazolyl, thiazolyl,  
 5 cyclohexyl, naphthyl, cyclohexenyl, pyrazolyl, benzothienyl, indolyl, 1,1,3-trioxo-2,3-  
 dihydro-1,2-benzisothiazolyl, 1,3-benzodioxolyl, cyclopentyl, tetrahydropyranyl, 1-  
 oxooctahydropyrido[1,2-a]pyrazinyl, 1,2,3,4-tetrahydronaphthyl, piperidinyl and  
 benzthiazolyl.

Ring A is pyrid-2-yl, pyrid-3-yl, phenyl, thien-2-yl, fur-2-yl, pyrazin-2-yl, 1,2,3-  
 10 thiadiazol-5-yl, thiazol-2-yl, thiazol-5-yl, cyclohexyl, naphtha-2-yl, cyclohex-1-enyl, pyrazol-  
 3-yl, benzothien-2-yl, indol-5-yl, 1,1,3-trioxo-2,3-dihydro-1,2-benzisothiazol-6-yl, 1,3-  
 benzodioxol-5-yl, cyclopentyl, tetrahydropyran-4-yl, 1-oxooctahydropyrido[1,2-a]pyrazin-7-  
 yl, 1,2,3,4-tetrahydronaphth-2-yl, piperidin-4-yl and benzthiazol-2-yl.

$R^1$  is selected from halo, cyano,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{1-4}$ alkoxy or  $C_{1-4}$ alkanoyl.

15  $R^1$  is selected from fluoro, chloro, cyano, methyl, 1-propenyl, methoxy or acetyl.

$R^1$  is selected from halo, nitro, cyano, sulphamoyl,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  
 $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl, tri- $(C_{1-4}$ alkyl)silyloxy, carbocyclyl and heterocyclyl $C_{0-4}$ alkylene-Y-;  
 wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^6$ .

20  $R^1$  is selected from fluoro, chloro, bromo, iodo, nitro, cyano, sulphamoyl, methyl,  
 ethyl, *t*-butyl, allyl, ethynyl, methoxy, isopropoxy, acetyl, dimethyl-*t*-butylsilyloxy, phenyl  
 and pyrimidin-4-ylamino; wherein  $R^1$  may be optionally substituted on carbon by one or more  
 groups selected from  $R^6$ .

$R^1$  is selected from halo, nitro, cyano, sulphamoyl,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  
 $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl, tri- $(C_{1-4}$ alkyl)silyloxy, carbocyclyl and heterocyclyl $C_{0-4}$ alkylene-Y-;  
 25 wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^6$ ;  
 wherein

Y is  $-NR^{12}$ ;

$R^{12}$  is hydrogen; and

$R^6$  is selected from halo,  $C_{2-4}$ alkenyl,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoylamino and  
 30 carbocyclyl.

$R^1$  is selected from fluoro, chloro, bromo, iodo, nitro, cyano, sulphamoyl, methyl,  
 ethyl, *t*-butyl, allyl, ethynyl, methoxy, isopropoxy, acetyl, dimethyl-*t*-butylsilyloxy, phenyl

and pyrimidin-4-ylamino; wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^6$ ; wherein

$R^6$  is selected from fluoro, chloro, ethenyl, acetyl, acetylamino and phenyl.

$R^1$  is selected from fluoro, chloro, bromo, iodo, nitro, cyano, sulphamoyl, methyl,

- 5 allyl, *t*-butyl, ethynyl, methoxy, isopropoxy, acetyl, allyloxy, trifluoromethyl, phenyl, benzyloxy, 4-chlorophenyl, 3-oxobutyl, 2-chloropyrimidin-4-yl, acetamidomethyl and dimethyl-*t*-butylsilyloxy.

Y is  $-NR^{12}$ .

$R^{12}$  is hydrogen.

- 10  $R^6$  is selected from halo,  $C_{2-4}$ alkenyl,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoylamino and carbocyclyl.

$R^6$  is selected from fluoro, chloro, ethenyl, acetyl, acetylamino and phenyl.

When Ring A is phenyl,  $R^1$  is selected from 2-fluoro, 3-fluoro, 4-fluoro, 2,4-difluoro, 3-chloro, 3-cyano, 4-cyano, 3-methyl, 3-(1-propenyl), 3-methoxy, 4-methoxy or 4-acetyl.

- 15 n is 0-2; wherein the values of  $R^1$  may be the same or different.

n is 0.

n is 1.

n is 2.

n is 3.

- 20  $R^1$ , n and Ring A together form phenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 3-chlorophenyl, 4-chlorophenyl, 3-bromophenyl, 4-bromophenyl, 3-iodophenyl, 4-iodophenyl, 3-methylphenyl, 4-*t*-butylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3-isopropoxyphenyl, 4-isopropoxyphenyl, 3-cyanophenyl, 4-cyanophenyl, 4-trifluoromethylphenyl, 4-sulphamoylphenyl, 3-nitrophenyl, 4-nitrophenyl, 3-acetylphenyl, 4-acetylphenyl, 3-allylphenyl, 3-allyloxyphenyl, 4-allyloxyphenyl, 4-ethynylphenyl, 3-benzyloxyphenyl, 4-benzyloxyphenyl, 4-(3-oxobutyl)phenyl, 4-(dimethyl-*t*-butylsilyloxy)phenyl, 4-(2-chloropyrimidin-4-ylamino)phenyl, 4-(acetamidomethyl)phenyl, 2,4-difluorophenyl, 3,5-dimethylphenyl, 3,5-dibenzyloxyphenyl, 3-methoxy-4-chlorophenyl, 3-fluoro-4-chlorophenyl, 3-cyano-4-methoxyphenyl, 3-iodo-4-methoxyphenyl, 3-nitro-4-chlorophenyl, 3,4,5-trimethoxyphenyl, biphenyl-3-yl, biphenyl-4-yl, cyclohexyl, 6-cyanonaphth-2-yl, cyclohex-1-en-1-yl, cyclopentyl, 3-phenylcyclopentyl, pyrid-2-yl, 2-methylpyrid-5-yl, thien-2-yl, 5-chlorothien-2-yl, 3-chloro-4-methylthien-2-yl, fur-2-yl, pyrazin-2-yl, 1,2,3-thiadiazol-5-yl, thiazol-2-yl, thiazol-5-yl, 4,5-dichlorothiazol-3-yl, 5-(4-

chlorophenyl)pyrazol-3-yl, benzothien-2-yl, indol-5-yl, 1,1,3-trioxo-2,3-dihydro-1,2-benzisothiazol-6-yl, 1,3-benzodioxol-5-yl, tetrahydropyran-4-yl, 1-oxooctahydropyrido[1,2-a]pyrazin-7-yl, 1,2,3,4-tetrahydronaphth-2-yl, 1-(pyrid-4-yl)piperidin-4-yl and benzothiazol-2-yl.

- 5         $R^2$  and  $R^3$  are independently selected from hydrogen or  $C_{1-4}$ alkyl.  
          $R^2$  and  $R^3$  are independently selected from hydrogen or methyl.  
          $R^2$  and  $R^3$  are both hydrogen.  
          $R^2$  and  $R^3$  are both methyl.  
         One of  $R^2$  and  $R^3$  is hydrogen and the other is methyl.
- 10         $R^4$  and  $R^5$  are independently selected from  $C_{1-4}$ alkyl; wherein  $R^4$  and  $R^5$  may be optionally substituted on carbon by one or more groups selected from  $R^{10}$ ; and  
          $R^{10}$  is selected from  $C_{1-4}$ alkoxy and  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino.  
         one of  $R^4$  and  $R^5$  is selected from hydrogen and  $C_{1-4}$ alkyl and the other is selected from  $C_{1-4}$ alkyl; wherein  $R^4$  and  $R^5$  may be optionally substituted on carbon by one or more
- 15        groups selected from  $R^{10}$ ; and  
          $R^{10}$  is selected from  $C_{1-4}$ alkoxy and  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino.  
         one of  $R^4$  and  $R^5$  is selected from selected from hydrogen, methyl, isopropyl and ethyl, and the other is selected from methyl, isopropyl, propyl and ethyl; wherein  $R^4$  and  $R^5$  may be optionally substituted on carbon by one or more groups selected from  $R^{10}$ .
- 20         $R^4$  and  $R^5$  are independently selected from methyl, ethyl, propyl and isopropyl; wherein  $R^4$  and  $R^5$  may be optionally substituted on carbon by one or more groups selected from  $R^{10}$ ; and  
          $R^{10}$  is selected from methoxy and  $N,N$ -dimethylamino.  
         one of  $R^4$  and  $R^5$  is selected from selected from hydrogen, methyl, isopropyl and ethyl,
- 25        and the other is selected from methyl, isopropyl, propyl and ethyl; wherein  $R^4$  and  $R^5$  may be optionally substituted on carbon by one or more groups selected from  $R^{10}$ ; and  
          $R^{10}$  is selected from methoxy, isopropoxy and  $N,N$ -dimethylamino.  
          $R^4$  and  $R^5$  are independently selected from methyl, ethyl, 2-methoxyethyl, 2-( $N,N$ -dimethylamino)propyl and isopropyl.
- 30        one of  $R^4$  and  $R^5$  is selected from selected from hydrogen, methyl, isopropyl and ethyl; and the other is selected from methyl, isopropyl, propyl, 2-methoxyethyl, 2-dimethylaminoethyl, 2-(isopropoxy)ethyl and ethyl.  
          $R^{10}$  is selected from  $C_{1-4}$ alkoxy and  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino.



R<sup>10</sup> is selected from methoxy, isopropoxy and *N,N*-dimethylamino.

R<sup>4</sup> and R<sup>5</sup> together with the nitrogen to which they are attached form isopropylamino, dimethylamino, diethylamino, diisopropylamino, *N*-(methyl)-*N*-(propyl)amino, *N*-(methyl)-*N*-(isopropyl)amino, *N*-(methyl)-*N*-(2-methoxyethyl)amino, *N*-(isopropyl)-*N*-(2-methoxyethyl)amino, *N*-(isopropyl)-*N*-[2-(isopropoxy)ethyl]amino, *N*-(methyl)-*N*-(2-dimethylaminoethyl)amino and *N*-(ethyl)-*N*-(isopropyl)amino.

Therefore in a further aspect of the invention there is provided the use of a compound of formula (I) (as depicted above) wherein:

- Ring A is pyridyl, phenyl, thienyl, furyl or pyrazinyl;
- 10 R<sup>1</sup> is selected from halo, cyano, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>1-4</sub>alkoxy or C<sub>1-4</sub>alkanoyl; n is 0-2; wherein the values of R<sup>1</sup> may be the same or different;
- R<sup>2</sup> and R<sup>3</sup> are independently selected from hydrogen or C<sub>1-4</sub>alkyl;
- R<sup>4</sup> and R<sup>5</sup> are independently selected from C<sub>1-4</sub>alkyl; wherein R<sup>4</sup> and R<sup>5</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>10</sup>; and
- 15 R<sup>10</sup> is selected from C<sub>1-4</sub>alkoxy and *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino;
- or a pharmaceutically acceptable salt thereof;
- in the manufacture of a medicament for use in the inhibition of 11 $\beta$ HSD1.

Therefore in a further aspect of the invention there is provided the use of a compound of formula (I) (as depicted above) wherein:

- 20 Ring A is pyrid-2-yl, phenyl, thien-2-yl, fur-2-yl, pyrazin-2-yl;
- R<sup>1</sup> is selected from fluoro, chloro, cyano, methyl, 1-propenyl, methoxy or acetyl;
- n is 0-2; wherein the values of R<sup>1</sup> may be the same or different;
- R<sup>2</sup> and R<sup>3</sup> are independently selected from hydrogen or methyl;
- R<sup>4</sup> and R<sup>5</sup> are independently selected from methyl, ethyl, 2-methoxyethyl, 2-(*N,N*-
- 25 dimethylamino)propyl and isopropyl;
- or a pharmaceutically acceptable salt thereof;
- in the manufacture of a medicament for use in the inhibition of 11 $\beta$ HSD1.

Therefore in a further aspect of the invention there is provided the use of a compound of formula (I) (as depicted above) wherein:

- 30 Ring A is carbocyclyl or heterocyclyl;
- R<sup>1</sup> is selected from halo, nitro, cyano, sulphonamoyl, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, tri-(C<sub>1-4</sub>alkyl)silyloxy, carbocyclyl and heterocyclylC<sub>0-4</sub>alkylene-Y-;

wherein R<sup>1</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>6</sup>;  
wherein:

Y is -NR<sup>12</sup>-;

R<sup>12</sup> is hydrogen; and

5 R<sup>6</sup> is selected from halo, C<sub>2-4</sub>alkenyl, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoylamino and carbocyclyl;

n is 0-3; wherein the values of R<sup>1</sup> may be the same or different;

R<sup>2</sup> and R<sup>3</sup> are independently selected from hydrogen or C<sub>1-4</sub>alkyl;

one of R<sup>4</sup> and R<sup>5</sup> is selected from hydrogen and C<sub>1-4</sub>alkyl and the other is selected  
10 from C<sub>1-4</sub>alkyl; wherein R<sup>4</sup> and R<sup>5</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>10</sup>; and

R<sup>10</sup> is selected from C<sub>1-4</sub>alkoxy and *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino;

or a pharmaceutically acceptable salt thereof;

in the manufacture of a medicament for use in the inhibition of 11βHSD1.

15 Therefore in a further aspect of the invention there is provided the use of a compound of formula (I) (as depicted above) wherein:

R<sup>1</sup>, n and Ring A together form phenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 3-chlorophenyl, 4-chlorophenyl, 3-bromophenyl, 4-bromophenyl, 3-iodophenyl, 4-iodophenyl, 3-methylphenyl, 4-*t*-butylphenyl, 3-methoxyphenyl, 4-  
20 methoxyphenyl, 3-isopropoxyphenyl, 4-isopropoxyphenyl, 3-cyanophenyl, 4-cyanophenyl, 4-trifluoromethylphenyl, 4-sulphamoylphenyl, 3-nitrophenyl, 4-nitrophenyl, 3-acetylphenyl, 4-acetylphenyl, 3-allylphenyl, 3-allyloxyphenyl, 4-allyloxyphenyl, 4-ethynylphenyl, 3-benzyloxyphenyl, 4-benzyloxyphenyl, 4-(3-oxobutyl)phenyl, 4-(dimethyl-*t*-butylsilyloxy)phenyl, 4-(2-chloropyrimidin-4-ylamino)phenyl, 4-(acetamidomethyl)phenyl,  
25 2,4-difluorophenyl, 3,5-dimethylphenyl, 3,5-dibenzyloxyphenyl, 3-methoxy-4-chlorophenyl, 3-fluoro-4-chlorophenyl, 3-cyano-4-methoxyphenyl, 3-iodo-4-methoxyphenyl, 3-nitro-4-chlorophenyl, 3,4,5-trimethoxyphenyl, biphenyl-3-yl, biphenyl-4-yl, cyclohexyl, 6-cyanonaphth-2-yl, cyclohex-1-en-1-yl, cyclopentyl, 3-phenylcyclopentyl, pyrid-2-yl, 2-methylpyrid-5-yl, thien-2-yl, 5-chlorothien-2-yl, 3-chloro-4-methylthien-2-yl, fur-2-yl,  
30 pyrazin-2-yl, 1,2,3-thiadiazol-5-yl, thiazol-2-yl, thiazol-5-yl, 4,5-dichlorothiazol-3-yl, 5-(4-chlorophenyl)pyrazol-3-yl, benzothien-2-yl, indol-5-yl, 1,1,3-trioxo-2,3-dihydro-1,2-benzisothiazol-6-yl, 1,3-benzodioxol-5-yl, tetrahydropyran-4-yl, 1-oxooctahydropyrido[1,2-

a]pyrazin-7-yl, 1,2,3,4-tetrahydronaphth-2-yl, 1-(pyrid-4-yl)piperidin-4-yl and benzothiazol-2-yl;

$R^2$  and  $R^3$  are independently selected from hydrogen or methyl; and

$R^4$  and  $R^5$  together with the nitrogen to which they are attached form isopropylamino, dimethylamino, diethylamino, diisopropylamino, *N*-(methyl)-*N*-(propyl)amino, *N*-(methyl)-*N*-(isopropyl)amino, *N*-(methyl)-*N*-(2-methoxyethyl)amino, *N*-(isopropyl)-*N*-(2-methoxyethyl)amino, *N*-(isopropyl)-*N*-[2-(isopropoxy)ethyl]amino, *N*-(methyl)-*N*-(2-dimethylaminoethyl)amino and *N*-(ethyl)-*N*-(isopropyl)amino; or a pharmaceutically acceptable salt thereof;


10 in the manufacture of a medicament for use in the inhibition of 11 $\beta$ HSD1.

In another aspect of the invention, preferred compounds of the invention are any one of the Examples or a pharmaceutically acceptable salt thereof.

In another aspect of the invention, preferred compounds of the invention are any one of the Reference Examples or a pharmaceutically acceptable salt thereof.

15 Another aspect of the present invention provides a process for preparing a compound of formula (I) or (Ia) or a pharmaceutically acceptable salt thereof which process (wherein variable groups are, unless otherwise specified, as defined in formula (I) or (Ia)) comprises of:


*Process 1*): reacting a compound of formula (II):



20

(II)


wherein V is a displaceable group; with an organometallic reagent of formula (III):



(III)


25 wherein M is a metal reagent;

*Process 2*): reacting a compound of formula (IV):



(IV)


wherein L is a displaceable group; with an amine of formula (V):



(V)


5

Process 3): reacting a compound of formula (VI):



(VI)

wherein  $R^xOC(O)-$  is an ester with a compound of formula (VII):



(VII)

10

and thereafter if necessary or desirable:

- i) converting a compound of the formula (I) into another compound of the formula (I);
- ii) removing any protecting groups;
- 15 iii) forming a pharmaceutically acceptable salt thereof.

L is a displaceable group, suitable values for L include halo, particularly fluoro or chloro.

V is a displaceable group, suitable values for V include the Weinreb amide *N*-methyl-*N*-methoxyamine.

- 20 M is a metal reagent. Suitable values for M include Grignard reagents such as MgBr and lithium.

The group  $R^xOC(O)-$  is an ester. Suitable values for  $R^x$  are methyl and ethyl.

The reactions described above may be performed under standard conditions. The intermediates described above are commercially available, are known in the art or may be prepared by known procedures.

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogeno group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard practice (for illustration see T.W. Green, Protective Groups in Organic Synthesis, John Wiley and Sons, 1991). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali

metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

As stated hereinbefore the compounds defined in the present invention possess 11 $\beta$ HSD1 inhibitory activity. These properties may be assessed using the following assay.

#### Assay

HeLa cells (human cervical carcinoma derived cells) were stably transfected with a construct containing four copies of the glucocorticoid response element (GRE) linked to a beta-galactosidase reporter gene (3 kb lac Z gene derived from pSV-B-galactosidase). These cells were then further stably transfected with a construct containing full-length human 11 $\beta$ HSD1 enzyme (in pCMVHyg) to create GRE4- $\beta$ Gal/11 $\beta$ HSD1 cells. The principal of the assay is as follows. Cortisone is freely taken up by the cells and is converted to cortisol by

11 $\beta$ HSD1 oxo-reductase activity and cortisol (but not cortisone) binds to and activates the glucocorticoid receptor. Activated glucocorticoid receptor then binds to the GRE and initiates transcription and translation of  $\beta$ -galactosidase. Enzyme activity can then be assayed with high sensitivity by colourimetric assay. Inhibitors of 11 $\beta$ HSD1 will reduce the conversion of  
5 cortisone to cortisol and hence decrease the production of  $\beta$ -galactosidase.

Cells were routinely cultured in DMEM (Invitrogen, Paisley, Renfrewshire, UK) containing 10% foetal calf serum (LabTech), 1% glutamine (Invitrogen), 1% penicillin & streptomycin (Invitrogen), 0.5 mg/ml G418 (Invitrogen) & 0.5mg/ml hygromycin (Boehringer). Assay media was phenol red free-DMEM containing 1% glutamine, 1%  
10 penicillin & streptomycin.

Compounds (1mM) to be tested were dissolved in dimethyl sulphoxide (DMSO) and serially diluted into assay media containing 10% DMSO. Diluted compounds were then plated into transparent flat-bottomed 384 well plates (Matrix, Hudson NH, USA).

The assay was carried out in 384 well microtitre plate (Matrix) in a total volume of  
15 50 $\mu$ l assay media consisting of cortisone (Sigma, Poole, Dorset, UK, 1 $\mu$ M), HeLa GRE4- $\beta$ Gal/11 $\beta$ HSD1 cells (10,000 cells) plus test compounds (3000 to 0.01 nM). The plates were then incubated in 5% O<sub>2</sub>, 95% CO<sub>2</sub> at 37°C overnight.

The following day plates were assayed by measurement of  $\beta$ -galactosidase production.

A cocktail (25 $\mu$ l) consisting of 10X Z-buffer (600 mM Na<sub>2</sub>HPO<sub>4</sub>, 400 mM  
20 NaH<sub>2</sub>PO<sub>4</sub>.2H<sub>2</sub>O, 100 mM KCl, 10 mM MgSO<sub>4</sub>.7H<sub>2</sub>O, 500 mM  $\beta$ -mercaptoethanol, pH 7.0), SDS (0.2%), chlorophenol red- $\beta$ -D-galactopyranoside (5mM, Roche Diagnostics) was added per well and plates incubated at 37°C for 3-4hours.  $\beta$ -Galactosidase activity was indicated by a yellow to red colour change (absorbance at 570nm) measured using a Tecan Spectrafluor Ultra.

25 The calculation of median inhibitory concentration (IC<sub>50</sub>) values for the inhibitors was performed using Origin 6.0 (Microcal Software, Northampton MA USA). Dose response curves for each inhibitor were plotted as OD units at each inhibitor concentration with relation to a maximum signal (cortisone, no compound) and IC<sub>50</sub> values calculated. Compounds of the present invention typically show an IC<sub>50</sub> <10 $\mu$ M.

30 According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (Ia) or a pharmaceutically acceptable salt thereof, or a compound selected from Examples, or a pharmaceutically acceptable salt thereof, as defined hereinbefore in association with a pharmaceutically acceptable diluent or

carrier.

The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical

5 administration as an ointment or cream or for rectal administration as a suppository.

In general the above compositions may be prepared in a conventional manner using conventional excipients.

The compound of formula (I), or a pharmaceutically acceptable salt thereof, will normally be administered to a warm-blooded animal at a unit dose within the range 0.1 –  
10 50 mg/kg that normally provides a therapeutically-effective dose. A unit dose form such as a tablet or capsule will usually contain, for example 1-1000 mg of active ingredient. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

15 We have found that the compounds defined in the present invention, or a pharmaceutically acceptable salt thereof, are effective 11 $\beta$ HSD1 inhibitors, and accordingly have value in the treatment of disease states associated with metabolic syndrome.

It is to be understood that where the term "metabolic syndrome" is used herein, this relates to metabolic syndrome as defined in 1) and/or 2) or any other recognised definition of  
20 this syndrome. Synonyms for "metabolic syndrome" used in the art include Reaven's Syndrome, Insulin Resistance Syndrome and Syndrome X. It is to be understood that where the term "metabolic syndrome" is used herein it also refers to Reaven's Syndrome, Insulin Resistance Syndrome and Syndrome X.

According to a further aspect of the present invention there is provided a compound of  
25 the formula (Ia) or a pharmaceutically acceptable salt thereof, or a compound selected from Examples, or a pharmaceutically acceptable salt thereof, as defined hereinbefore for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

Thus according to this aspect of the invention there is provided a compound of the formula (Ia) or a pharmaceutically acceptable salt thereof, or a compound selected from  
30 Examples, or a pharmaceutically acceptable salt thereof, as defined hereinbefore for use as a medicament.

According to another feature of the invention there is provided the use of a compound of the formula (Ia) or a pharmaceutically acceptable salt thereof, or a compound selected from



the Examples, or a pharmaceutically acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an 11 $\beta$ HSD1 inhibitory effect in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound  
5 selected from Reference Examples, or a pharmaceutically acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an 11 $\beta$ HSD1 inhibitory effect in a warm-blooded animal, such as man.

Where production of or producing an 11 $\beta$ HSD1 inhibitory effect is referred to suitably this refers to the treatment of metabolic syndrome. Alternatively, where production of an  
10 11 $\beta$ HSD1 inhibitory effect is referred to this refers to the treatment of diabetes, obesity, hyperlipidaemia, hyperglycaemia, hyperinsulinemia or hypertension, particularly diabetes and obesity. Alternatively, where production of an 11 $\beta$ HSD1 inhibitory effect is referred to this refers to the treatment of glaucoma, osteoporosis, tuberculosis, dementia, cognitive disorders or depression.

15 According to a further feature of this aspect of the invention there is provided a method for producing an 11 $\beta$ HSD1 inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

According to a further feature of this aspect of the invention there is provided a  
20 method for producing an 11 $\beta$ HSD1 inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (Ia) or a pharmaceutically acceptable salt thereof, or a compound selected from Examples, or a pharmaceutically acceptable salt thereof.

According to a further feature of this aspect of the invention there is provided a  
25 method for producing an 11 $\beta$ HSD1 inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound selected from the Reference Examples, or a pharmaceutically acceptable salt thereof.

In addition to their use in therapeutic medicine, the compounds of formula (I), or a  
30 pharmaceutically acceptable salt thereof, are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of 11 $\beta$ HSD1 in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

The inhibition of 11 $\beta$ HSD1 described herein may be applied as a sole therapy or may involve, in addition to the subject of the present invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment.

5 Simultaneous treatment may be in a single tablet or in separate tablets. For example agents than might be co-administered with 11 $\beta$ HSD1 inhibitors, particularly those of the present invention, may include the following main categories of treatment:

- 1) Insulin and insulin analogues;
- 10 2) Insulin secretagogues including sulphonylureas (for example glibenclamide, glipizide) and prandial glucose regulators (for example repaglinide, nateglinide);
- 3) Insulin sensitising agents including PPAR $\gamma$  agonists (for example pioglitazone and rosiglitazone);
- 4) Agents that suppress hepatic glucose output (for example metformin);
- 15 5) Agents designed to reduce the absorption of glucose from the intestine (for example acarbose);
- 6) Agents designed to treat the complications of prolonged hyperglycaemia; e.g. aldose reductase inhibitors
- 7) Other anti-diabetic agents including phosphotyrosine phosphatase inhibitors, glucose 6 - phosphatase inhibitors, glucagon receptor antagonists, glucokinase activators,
- 20 glycogen phosphorylase inhibitors, fructose 1,6 bisphosphatase inhibitors, glutamine:fructose -6-phosphate amidotransferase inhibitors
- 8) Anti-obesity agents (for example sibutramine and orlistat);
- 9) Anti- dyslipidaemia agents such as, HMG-CoA reductase inhibitors (statins, eg pravastatin); PPAR $\alpha$  agonists (fibrates, eg gemfibrozil); bile acid sequestrants
- 25 (cholestyramine); cholesterol absorption inhibitors (plant stanols, synthetic inhibitors); ileal bile acid absorption inhibitors (IBATi), cholesterol ester transfer protein inhibitors and nicotinic acid and analogues (niacin and slow release formulations);
- 10) Antihypertensive agents such as,  $\beta$  blockers (eg atenolol, inderal); ACE inhibitors (eg lisinopril); calcium antagonists (eg. nifedipine); angiotensin receptor antagonists (eg
- 30 candesartan),  $\alpha$  antagonists and diuretic agents (eg. furosemide, benzthiazide);
- 11) Haemostasis modulators such as, antithrombotics, activators of fibrinolysis and antiplatelet agents; thrombin antagonists; factor Xa inhibitors; factor VIIa inhibitors);

antiplatelet agents (eg. aspirin, clopidogrel); anticoagulants (heparin and Low molecular weight analogues, hirudin) and warfarin; and

- 12) Anti-inflammatory agents, such as non-steroidal anti-inflammatory drugs (eg. aspirin) and steroidal anti-inflammatory agents (eg. cortisone).

5 In the above other pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

### Examples

The invention will now be illustrated in the following non limiting Examples, in which  
10 standard techniques known to the skilled chemist and techniques analogous to those described in these Examples may be used where appropriate, and in which, unless otherwise stated:  
(i) evaporations were carried out by rotary evaporation in vacuo and work up procedures were carried out after removal of residual solids such as drying agents by filtration;  
(ii) all reactions were carried out under an inert atmosphere at ambient temperature, typically  
15 in the range 18-25°C, with solvents of HPLC grade under anhydrous conditions, unless otherwise stated;  
(iii) column chromatography (by the flash procedure) was performed on Silica gel 40-63 µm (Merck);  
(iv) yields are given for illustration only and are not necessarily the maximum attainable;  
20 (v) the structures of the end products of the formula (I) were generally confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; magnetic resonance chemical shift values were measured in deuterated CDCl<sub>3</sub> (unless otherwise stated) on the delta scale (ppm downfield from tetramethylsilane); proton data is quoted unless otherwise stated; spectra were recorded on a Varian Mercury-300 MHz, Varian Unity plus-  
25 400 MHz, Varian Unity plus-600 MHz or on Varian Inova-500 MHz spectrometer unless otherwise stated data was recorded at 400MHz; and peak multiplicities are shown as follows:  
s, singlet; d, doublet; dd, double doublet; t, triplet; tt, triple triplet; q, quartet; tq, triple quartet; m, multiplet; br, broad; ABq, AB quartet; ABd, AB doublet, ABdd, AB doublet of doublets; dABq, doublet of AB quartets; LCMS were recorded on a Waters ZMD, LC column xTerra  
30 MS C<sub>8</sub>(Waters), detection with a HP 1100 MS-detector diode array equipped; mass spectra (MS) (loop) were recorded on VG Platform II (Fisons Instruments) with a HP-1100 MS-detector diode array equipped; unless otherwise stated the mass ion quoted is (MH<sup>+</sup>);

unless further details are specified in the text, analytical high performance liquid chromatography (HPLC) was performed on Prep LC 2000 (Waters), Cromasil C<sub>8</sub>, 7 µm, (Akzo Nobel); MeCN and de-ionised water 10 mM ammonium acetate as mobile phases, with suitable composition;

- 5 (vii) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), HPLC, infra-red (IR), MS or NMR analysis;  
(viii) where solutions were dried magnesium sulphate was the drying agent;  
(ix) where an "ISOLUTE" column is referred to, this means a column containing 2 g of silica, the silica being contained in a 6 ml disposable syringe and supported by a porous disc of 54Å pore size, obtained from International Sorbent Technology under the name "ISOLUTE";  
10 "ISOLUTE" is a registered trade mark;

(x) the following abbreviations may be used hereinbefore or hereinafter:-

DCM dichloromethane;

EtOAc ethyl acetate;

15 MeCN acetonitrile; and

THF tetrahydrofuran.

### Reference Example 1

#### (N,N-Dimethylsulphamoylmethyl)(phenyl)ketone

- 20 The title compound was prepared by the procedure of J.Med.Chem.; EN; 30; 12; 1987; 2232-2239. Reference Example 1 is exemplified in this reference.

### Reference Example 2

#### (N,N-Dimethylsulphamoylmethyl)(4-fluorophenyl)ketone

- 25 The title compound was prepared by the procedure of Reference Example 1. NMR: 2.9 (s, 6H), 4.5 (s, 2H), 7.2 (m, 2H), 8.0 (m, 2H); m/z 244.

### Reference Example 3

#### (N,N-Dimethylsulphamoylmethyl)(thien-2-yl)ketone

- 30 To a stirred solution of methylthiophene-2-carboxylate (520mg, 3.65mmol) and N,N-dimethylmethanesulphonamide (375mg, 3.04mmol) in ethylene glycol dimethyl ether (15ml) was added sodium hydride (60% suspension in oil, 328mg, 8.21mmol). The reaction was warmed to 85°C and stirred at this temperature overnight then cooled to room temperature and
-

quenched with water. The resulting brown solution was acidified to ~pH2 with concentrated hydrochloric acid and then extracted with DCM (2 x 40ml). The organic layers were combined, washed with water (30ml) and brine (20ml) then dried, filtered and evaporated to yield crude product. This was purified by column chromatography (20g Silica, eluting with DCM) to yield an oil which crystallised on standing. This material was still impure. The crude was product partitioned between DCM and 1M sodium hydroxide solution, the layers separated and the sodium hydroxide layer re extracted with DCM. The aqueous layer was then acidified to ~pH3 with concentrated HCl and then extracted with DCM twice. These two DCM layers were combined, washed with brine, dried, filtered and evaporated to yield the product as a solid (56mg, 7%). NMR: 2.90 (s, 6H), 4.45 (s, 2H), 7.20 (m, 1H), 7.75 (m, 1H), 7.90 (m, 1H); m/z: 234.

#### Reference Example 4

##### [1-(*N,N*-Dimethylsulphamoyl)ethyl](phenyl)ketone

To a stirred solution of (*N,N*-dimethylsulphamoylmethyl)(phenyl)ketone (Reference Example 1; 88mg, 0.39mmol) in DMF (7ml) was added potassium carbonate (107mg, 0.78mmol) followed by methyl iodide (113mg, 0.8mmol). The resulting suspension was stirred at room temperature for 2 hours. The reaction was quenched with water (~50ml) and extracted with DCM (2x50ml). The organic layers combined, dried, filtered and evaporated to yield the product as a yellow oil (still contains a trace of DMF). NMR: 1.70 (d, 3H), 2.90 (s, 6H), 5.15 (m, 1H), 7.50 (t, 2H), 7.60 (m, 2H), 8.00 (br m, 1H); m/z: 242.

#### Example 1

##### [1-(*N,N*-Dimethylsulphamoyl)-1-methylethyl](phenyl)ketone

To a stirred solution of [1-(*N,N*-dimethylsulphamoyl)ethyl](phenyl)ketone (Reference Example 4; 33mg, 0.14mmol) in DMF was added potassium carbonate (39mg, 0.28mmol) and methyl iodide (60mg, 0.42mmol). The reaction was warmed to 40°C and stirred at this temperature for 18 hours. Further methyl iodide was added (60mg, 0.42mmol) and the reaction was stirred at 40°C for a further 24 hours. The volatiles were removed under reduced pressure and the resulting crude product was partitioned between ether and 1M sodium hydroxide solution, the ether layer was separated and re-extracted with sodium hydroxide solution then washed with brine, dried, filtered and evaporated to yield the product as a clear oil (16mg, 43%). M/z: 256.

**Example 2**(*N,N*-Dimethylsulphamoylmethyl)(pyrid-2-yl)ketone

To a solution of *N*-dimethylmethanesulphonamide (1.23 g, 10mmol) in THF (30ml), under an inert atmosphere at 0°C was added dropwise a solution of butyl lithium in hexanes 5 1.6M (12.5ml, 20mmol). After 30 minutes at 0°C, the mixture (white paste) was cooled to -78°C and ethylpicolinate (1.51g, 10mmol) in THF (5 ml) was added. After 1 hour, the cooling bath was removed and the temperature was allowed to warm to 0°C. The mixture was diluted with cooled water and extracted twice with ether. The aqueous phase was acidified to pH 5 and extracted three times with ethyl acetate. The ethyl acetate extracts were combined 10 and washed with brine, dried over, filtered and concentrated. The brown mauve resulting oil was triturated in ether until crystallisation occurred and the solid was filtered off (1.4 g ; 66%). NMR: 2.92 (s, 6H), 4.98 (s, 2H), 7.55 (m, 1H), 7.89 (m, 1H), 8.11 (d, 1H), 8.74 (m, 1H); m/z: 229.


**15 Example 3**(*N,N*-Diisopropylsulphamoylmethyl)(4-fluorophenyl)ketone

To a stirred solution of *N,N*-diisopropylmethanesulphonamide (120mg, 0.67mmol) in anhydrous THF (3ml) at -20°C was added a 1M solution of lithium bis(trimethylsilyl)amide in THF (1.34ml, 1.34mmol). The reaction was stirred at -20°C for 30 minutes and then a 20 solution of methyl-4-fluorobenzoate (134mg, 0.87mmol) in anhydrous THF (1ml) was added. The reaction was allowed to warm to room temperature over an hour then quenched with saturated ammonium chloride solution (5ml). The layers were separated and the aqueous layer was extracted with EtOAc. The THF and ethyl acetate extracts were combined, washed with brine, dried, filtered and evaporated to yield an impure oil. The crude product was purified by 25 column chromatography (eluting with DCM to 5%MeOH/DCM) to yield the product as an oil which crystallised on standing (70mg, 35%). NMR: 1.25 (d, 12H), 3.65 (m, 2H), 4.40 (s, 2H), 7.10 (t, 2H), 8.02 (m, 2H); m/z: 300 (M-H).

**Examples 4-75 and Reference Examples 5 and 6**

30 The procedure described in Example 3 was repeated using the appropriate reagent(s) in place of *N,N*-diisopropylmethanesulphonamide and/or methyl-4-fluorobenzoate to give the following Examples. Where the methanesulphonamides were not known compounds or commercially available the preparation of the starting materials (SM) is indicated.

---



Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
4	3-Chloro phenyl	Me	Me	2.95 (s, 6H), 4.55 (s, 2H), 7.45 (t, 1H), 7.60 (m, 1H), 7.90 (m, 1H), 8.00 (m, 1H)	262	
5	4-Fluoro phenyl	Et	Et	1.20 (t, 6H), 3.30 (q, 4H), 4.50 (s, 2H), 7.15 (t, 2H), 8.10 (m, 2H)	274	<sup>2</sup>
6	3-Methoxy phenyl	Me	Me	2.90 (s, 6H), 3.85 (s, 3H), 4.55 (s, 2H), 7.20 (m, 1H), 7.40 (t, 1H), 7.50 (m, 1H), 7.65 (m, 1H)	258	
7	Fur-2-yl	i-Pr	i-Pr	1.30 (d, 12H), 3.75 (m, 2H), 4.40 (s, 2H), 6.60 (m, 1H), 7.40 (m, 1H), 7.0 (m, 1H)	272 (M- H) <sup>+</sup>	<sup>3</sup>
8	4-Fluoro phenyl	Me	-(CH <sub>2</sub> ) <sub>2</sub> - OCH <sub>3</sub>	2.95 (s, 3H), 3.25 (s, 3H), 3.35 (q, 2H), 3.44 (q, 2H), 4.55 (s, 2H), 7.10 (t, 2H), 8.00 (m, 2H)	290	Meth 1
9	4-Fluoro phenyl	Me	Pr	0.90 (t, 3H), 1.60 (m, 2H), 2.90 (s, 3H), 3.15 (t, 2H), 4.55 (s, 2H), 7.15 (t, 2H), 8.10 (m, 2H)	274	Meth 2
10	4-Fluoro phenyl	Et	i-Pr	1.20 (t, 3H), 1.25 (d, 6H), 3.20 (q, 2H), 3.95 (m, 1H), 4.50 (s, 2H), 7.20 (t, 2H), 8.10 (m, 2H)	286 (M- H) <sup>+</sup>	Meth 3

Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
11	4-Methoxy phenyl	Me	Me	2.90 (s, 6H), 3.90 (s, 3H), 3.55 (s, 2H); 7.00 (d, 2H), 8.00 (d, 2H)	258	
12	Thiazol-2-yl	Me	Me	2.90 (s, 6H), 4.85 (s, 2H), 7.80 (d, 1H), 8.10 (d, 1H)	233 (M- H) <sup>+</sup>	
13	1,2,3-Thia- diazol-5-yl	Me	Me	2.95 (s, 6H), 4.50 (s, 2H), 9.20 (s, 1H)	234 (M- H) <sup>+</sup>	
14	Pyrazin-2-yl	Me	Me	2.95 (s, 6H), 4.80 (s, 2H), 8.70 (m, 1H), 8.80 (d, 1H), 9.30 (s, 1H)	228 (M- H) <sup>+</sup>	
15	4-Fluoro phenyl	Me	-(CH <sub>2</sub> ) <sub>2</sub> - N(Me) <sub>2</sub>	2.30 (s, 6H), 2.50 (t, 2H), 3.00 (br s, 3H), 3.35 (t, 2H), 4.70 (br s, 2H), 7.20 (br t; 3H), 8.10 (m, 2H)	303	Meth 6
16	Thiazol-5-yl	Me	Me	2.95 (s, 6H), 4.50 (s, 2H), 8.60 (s, 1H), 9.10 (s, 1H)	235	<sup>1</sup>
17 <sup>4</sup> , <sup>5</sup>	4-Trifluoro- methyl phenyl	Me	Me	2.9 (s, 6H), 4.6 (s, 2H), 7.8 (d, 2H), 8.2 (d, 2H)	294 (M- H) <sup>+</sup>	
18 <sup>4</sup> , <sup>5</sup>	4- <i>t</i> - Butylphenyl	Me	Me	1.6 (s, 9H), 2.9 (s, 6H), 4.6 (s, 2H), 7.6 (d, 2H), 8.0 (d, 2H)	282 (M- H) <sup>+</sup>	
19 <sup>4</sup>	6-Cyano naphth-2-yl	Me	Me		301 (M- H) <sup>+</sup>	
20 <sup>4</sup> , <sup>5</sup>	4-Chloro-3- fluorophenyl	Me	Me	2.9 (s, 6H), 4.6 (s, 2H), 7.6 (m, 1H), 7.8 (m, 2H)	278 (M- H) <sup>+</sup>	
21 <sup>4</sup>	3-Cyano-4- methoxy phenyl	Me	Me	2.9 (s, 6H), 4.1 (s, 3H), 4.5 (s, 2H), 7.1 (d, 1H), 8.3 (m, 2H)		



Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
22 <sup>4,5</sup>	3-Iodo phenyl	Me	Me	2.9 (s, 6H), 4.5 (s, 2H), 7.3 (dd, 1H), 8.0 (m, 2H), 8.4 (d, 1H)	352 (M-H) <sup>-</sup>	
23 <sup>4,5</sup>	3-Iodo-4-methoxy phenyl	Me	Me	2.9 (s, 6H), 4.0 (s, 3H), 4.5 (s, 2H), 6.9 (d, 1H), 8.0 (dd, 1H), 8.4 (d, 2H)	382 (M-H) <sup>-</sup>	
24 <sup>4,5</sup>	3-Bromo phenyl	Me	Me	2.9 (s, 6H), 4.5 (s, 2H), 7.4 (t, 1H), 7.7 (dd, 1H), 8.0 (dd, 1H), 8.2 (d, 1H)	306 (M-H) <sup>-</sup>	
25 <sup>4,5</sup>	Biphen-4-yl	Me	Me		302 (M-H) <sup>-</sup>	
26 <sup>4,5</sup>	3,5-Dimethyl phenyl	Me	Me		254 (M-H) <sup>-</sup>	
27 <sup>4,5</sup>	3,5-Dibenzoyloxy phenyl	Me	Me		438 (M-H) <sup>-</sup>	
RE 5 <sup>4,5</sup>	4-Nitro phenyl	Me	Me		271 (M-H) <sup>-</sup>	
28 <sup>4,5</sup>	3-Nitro phenyl	Me	Me		271 (M-H) <sup>-</sup>	
29 <sup>4,5</sup>	3-Nitro-4-chlorophenyl	Me	Me		305 (M-H) <sup>-</sup>	
30 <sup>4,5</sup>	6-Methyl pyrid-3-yl	Me	Me		241 (M-H) <sup>-</sup>	
31 <sup>6</sup>	2,4-Difluoro phenyl	i-Pr	i-Pr	d <sup>6</sup> -DMSO, 1.22 (d, 12H), 3.75 (m, 2H), 4.65 (s, 2H), 7.24 (m, 1H), 7.42 (m, 1H), 8.00 (m, 1H)	318 (M-H) <sup>-</sup>	<sup>3</sup>

Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
32 <sup>6</sup>	4-Bromo phenyl	Me	Me	d <sup>6</sup> -DMSO, 2.82 (s, 6H), 4.92 (s, 2H), 7.77 (d, 2H), 7.96 (d, 2H)	304 (M- H) <sup>+</sup>	
33 <sup>6</sup>	4-Iodo phenyl	Me	Me	d <sup>6</sup> -DMSO, 2.80 (s, 6H), 4.88 (s, 2H), 7.78 (d, 2H), 7.95 (d, 2H)	354	
34 <sup>6</sup>	3-(Allyloxy) phenyl	Me	Me	d <sup>6</sup> -DMSO, 2.81 (s, 6H), 4.63 (m, 2H), 4.90 (s, 2H), 5.27 (m, 1H), 5.41 (m, 1H), 6.05 (m, 1H), 7.27 (m, 1H), 7.46 (m, 1H), 7.54 (m, 1H), 7.63 (m, 1H)	284	
35 <sup>4,7</sup>	3-Cyano phenyl	i-Pr	i-Pr		307 (M- H) <sup>+</sup>	<sup>3</sup>
36 <sup>4,7</sup>	Phenyl	i-Pr	i-Pr		282 (M- H) <sup>+</sup>	<sup>3</sup>
37 <sup>4,6</sup>	1,3-Benzo- dioxol-5-yl	i-Pr	i-Pr	d <sup>6</sup> -DMSO, 1.22 (d, 12H), 3.74 (m, 2H), 4.66 (s, 2H), 6.15 (s, 2H), 7.04 (d, 1H), 7.46 (d, 1H), 7.66 (dd, 1H)	326 (M- H) <sup>+</sup>	<sup>3</sup>
38 <sup>4,7</sup>	4-Cyano phenyl	i-Pr	i-Pr		307 (M- H) <sup>+</sup>	<sup>3</sup>
39 <sup>4,6</sup>	4- (Acetamido- methyl) phenyl	i-Pr	i-Pr		353 (M- H) <sup>+</sup>	<sup>3</sup>
40 <sup>4,7</sup>	1,1,3- Trioxo-2,3- dihydro-1,2- benziso- thiazol-6-yl	i-Pr	i-Pr		387 (M- H) <sup>+</sup>	<sup>3,9</sup>

Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
41 <sup>4,7</sup>	1 <i>H</i> -indol-5-yl	i-Pr	i-Pr	d <sup>6</sup> -DMSO, 1.23 (d, 12H), 3.76 (m, 2H), 4.72 (s, 2H), 6.63 (m, 1H), 7.14 (br s, 1H), 7.47 (m, 2H), 7.76 (m, 1H), 8.38 (m, 1H)	321 (M-H) <sup>-</sup>	<sup>3</sup>
42 <sup>4,7</sup>	4-(Benzyloxy)phenyl	Me	Me	d <sup>6</sup> -DMSO, 2.79 (s, 6H), 4.81 (s, 2H), 5.23 (s, 2H), 7.15 (d, 2H), 7.41 (m, 5H), 8.00 (d, 2H)	334	
43 <sup>4,6</sup>	Biphen-3-yl	Me	Me		304	
44 <sup>4,6</sup>	3-Acetylphenyl	Me	Me	d <sup>6</sup> -DMSO, 2.64 (s, 3H), 2.83 (s, 6H), 5.00 (s, 2H), 7.71 (m, 1H), 8.24 (m, 2H), 8.50 (m, 1H)	270	
45 <sup>4,6</sup>	3-(Benzyloxy)phenyl	Me	Me		332 (M-H) <sup>-</sup>	
46 <sup>4,6</sup>	4,5-Dichlorothiazol-2-yl	Me	Me	d <sup>6</sup> -DMSO, 2.83 (s, 6H), 4.88 (s, 2H)	303 (M-H) <sup>-</sup>	
47 <sup>4,6</sup>	Benzothien-2-yl	Me	Me	d <sup>6</sup> -DMSO, 2.84 (s, 6H), 4.95 (s, 2H), 7.53 (m, 2H), 8.06 (m, 2H), 8.57 (s, 1H)	282 (M-H) <sup>-</sup>	
48 <sup>4,6</sup>	2-Chlorothien-5-yl	Me	Me		266 (M-H) <sup>-</sup>	
49 <sup>4,6</sup>	3-Chloro-4-methylthien-2-yl	Me	Me	d <sup>6</sup> -DMSO, 2.21 (s, 3H), 2.84 (s, 6H), 4.79 (s, 2H), 7.92 (s, 1H)	280 (M-H) <sup>-</sup>	
50 <sup>4,6</sup>	5-(4-Chlorophenyl)pyrazol-3-yl	Me	Me		326 (M-H) <sup>-</sup>	<sup>10</sup>

Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
51 <sup>4,5</sup>	4-(3-Oxobutyl)phenyl	Me	Me		296	
52 <sup>4,5</sup>	4-(2-Chloro pyrimidin-4-ylamino)phenyl	Me	Me	d <sup>6</sup> -DMSO, 2.81 (s, 6H), 4.82 (s, 2H), 6.90 (d, 1H), 7.81 (d, 2H), 8.05 (d, 2H), 8.27 (d, 2H), 10.48 (br s, 1H)	355	
53 <sup>4,5</sup>	4-Sulphamoylphenyl	Me	Me	d <sup>6</sup> -DMSO, 2.81 (s, 6H), 4.98 (s, 2H), 7.57 (s, 2H), 7.96 (d, 2H), 8.22 (d, 2H)	305 (M-H) <sup>-</sup>	
54 <sup>4,8</sup>	Benzothien-2-yl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> -O-iPr		382 (M-H) <sup>-</sup>	Meth 11
55 <sup>4,8</sup>	4-Fluorophenyl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> -O-iPr		344 (M-H) <sup>-</sup>	Meth 11
56 <sup>4,8</sup>	2,4-Difluorophenyl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> -O-iPr		362 (M-H) <sup>-</sup>	Meth 11
57 <sup>4,8</sup>	2-Chloro thien-5-yl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> -O-iPr		366 (M-H) <sup>-</sup>	Meth 11
58 <sup>4,8</sup>	Thiazol-2-yl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> -O-iPr		333 (M-H) <sup>-</sup>	Meth 11
59 <sup>4,8</sup>	4-Fluorophenyl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> OCH <sub>3</sub>		316 (M-H) <sup>-</sup>	Meth 10
60 <sup>4,8</sup>	2-Chloro thien-5-yl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> OCH <sub>3</sub>		338 (M-H) <sup>-</sup>	Meth 10
61 <sup>4,8</sup>	2,4-Difluorophenyl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> OCH <sub>3</sub>		334 (M-H) <sup>-</sup>	Meth 10
62	4-( <i>t</i> -Butyl dimethyl silyloxy)phenyl	i-Pr	i-Pr	0.27 (s, 6H), 1.00 (s, 9H), 1.33 (d, 12H), 3.73 (m, 2H), 4.47 (s, 2H), 6.90 (d, 2H), 7.99 (d, 2H)		11

Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
RE 6 <sup>4,5</sup>	Cyclohexyl	Me	Me	1.3 (m, 5H), 1.7 (d, 1H), 1.8 (b, 2H), 1.9 (b, 2H), 2.7 (m, 1H), 2.9 (s, 6H), 4.0 (s, 2H)		
63 <sup>4</sup>	Cyclohex-1- en-1-yl	Me	Me		232 (M+H) <sup>+</sup>	
64 <sup>7</sup>	3-Phenyl cyclopentyl	Me	Me	d <sup>6</sup> -DMSO, 1.80 (m, 6H), 2.27 (m, 1H), 2.80 (s, 6H), 3.04 (m, 1H), 4.37 (s, 2H), 7.22 (m, 5H)		
65 <sup>7</sup>	1,2,3,4- Tetrahydro naphth-2-yl	Me	Me	d <sup>6</sup> -DMSO, 1.62 (m, 1H), 2.14 (m, 1H), 2.80 (s, 6H), 2.93 (m, 5H), 4.50 (m, 2H), 7.07 (m, 4H)		
66 <sup>7</sup>	Cyclopentyl	Me	Me	d <sup>6</sup> -DMSO, 1.54 (m, 4H), 1.66 (m, 2H), 1.80 (m, 2H), 2.77 (s, 6H), 3.10 (m, 1H), 4.34 (s, 2H)		
67 <sup>7</sup>	1-(Pyrid-4- yl)- piperidin-4- yl	Me	Me	d <sup>6</sup> -DMSO, 1.43 (m, 2H), 1.92 (m, 2H), 2.78 (s, 6H), 2.90 (m, 3H), 3.92 (m, 2H), 4.42 (s, 2H), 6.80 (m, 2H), 8.12 (m, 2H)		
68 <sup>7</sup>	4-Tetrahyro pyran-4-yl	Me	Me	d <sup>6</sup> -DMSO, 1.43 (m, 2H), 1.77 (m, 2H), 2.77 (s, 6H), 2.86 (m, 1H), 3.33 (m, 2H), 3.85 (m, 2H), 4.39 (s, 2H)		

Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z <sup>-</sup>	SM
69 <sup>7</sup>	1-Oxo octahydro pyrido[1,2-a] pyrazin-7-yl	Me	Me	d <sup>6</sup> -DMSO, 1.23 (m, 2H), 2.05 (m, 2H), 2.16 (m, 1H), 2.37 (m, 2H), 2.76 (s, 6H), 2.86 (m, 2H), 3.05 (m, 2H), 3.22 (m, 1H), 4.40 (m, 2H), 7.60 (br s, 1H)		
70	Benzothien- 2-yl	i-Pr	i-Pr	1.30-1.23 (d, 6H); 3.68- 3.84 (m, 2H); 4.56 (s, 2H); 7.37-7.53 (m, 2H); 7.83- 7.90 (d, 1H); 7.91-7.99 (d, 1H); 8.22 (s, 1H)	+ve 340	<sup>3</sup>
71	Benzthiazol- 2-yl	i-Pr	i-Pr	1.26-1.45 (d, 12H); 3.74- 3.94 (m, 2H); 4.92 (s, 2H) 7.50 (m, 2H); 7.93-8.02 (d, 1H); 8.17-8.27 (d, 1H)	+ve 341	<sup>3</sup>
72	3- Isopropoxy phenyl	i-Pr	i-Pr	1.28-1.45 (m, 18H); 3.67- 3.83 (m, 2H); 4.51 (s, 2H); 4.55-4.69 (m, 1H); 7.08- 7.16 (app dd, 1H); 7.33- 7.43 (app t, 1H), 7.55 (m, 1H); 7.59-7.65 (d, 1H)	+ve 342	<sup>3</sup>
73	Thiazol-2-yl	i-Pr	i-Pr	1.23-1.43 (d, 12H); 3.72- 3.91 3(m, 2H); 4.80 (s, 2H); 7.71-7.79 (d, 1H); (8.03-8.10 (d, 1H)	+ve 291	<sup>3</sup>
74	4-Bromo phenyl	i-Pr	i-Pr	1.09-1.32 (d, 12H); 3.66- 3.86 (m, 2H); 4.76 (s, 2H); 7.71-7.85 (d, 2H); 7.90- 8.03 (d, 2H)	-ve 360	<sup>3</sup>

Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
75	4-Isopropoxy phenyl	i-Pr	i-Pr	1.23-1.40 (d, 12H); 1.40-1.42 (d, 6H); 3.66-3.80 (m, 2H); 4.48 (s, 2H); 4.60-4.72(m, 1H); 6.89-6.96 (d, 2H); 7.97- 8.07 (d, 2H)	+ve 342	<sup>3</sup>

<sup>1</sup> Starting ester prepared according to Tetrahedron Lett.; EN; 25; 51; 1984; 5939-5942

<sup>2</sup> Sulphonamide preparation: J.Amer.Chem.Soc.; 76; 1954; 303

<sup>3</sup> Sulphonamide preparation: Tetrahedron; EN; 25; 1969; 181-189

<sup>4</sup> Reaction carried out at room temperature overnight

5 <sup>5</sup> Product crystallised from ethyl acetate / hexane in place of chromatography.

<sup>6</sup> Purification by chromatography (eluting with 25% EtOAc/isohexane to 50% EtOAc/isohexane)

<sup>7</sup> Product triturated with diethyl ether.

<sup>8</sup> Purification by chromatography (eluting with 10% EtOAc/isohexane to 20%

10 EtOAc/isohexane)

<sup>9</sup> Starting ester prepared according to Azerbaidzhanskii Khimicheskii Zhurnal; 1997; 1-4; 62-66.


<sup>10</sup> Starting ester prepared according to PCT Int. Appl.; 1997; 99pp.; WO9721682.

<sup>11</sup> Starting ester prepared according to J. Org. Chem.; 1991; 56(16); 4884-7.

15

### Examples 76-78

The procedure described in Example 3 was repeated using the appropriate reagent(s) in place of *N,N*-diisopropylmethanesulphonamide to give the following Examples.



Ex	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
76	Me	Me	1.73 (s, 6H), 2.93 (s, 6H), 7.10 (m, 2H), 8.05 (m, 2H)		Method 7
77	Me	i-Pr	1.12 (d, 6H), 1.62 (s, 6H), 2.68 (s, 3H), 3.87 (m, 1H), 7.30 (m, 2H), 7.94 (m, 2H)	302	Method 8

Ex	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
78	H	i-Pr	1.28 (d, 6H), 2.10 (s, 6H), 3.80 (m, 1H), 7.07 (m, 2H), 7.87 (m, 2H), 11.40 (br s, 1H)		Method 9

**Example 79**(*N,N*-Dimethylsulphamoylmethyl)(4-chlorophenyl)ketone

To a stirred solution of methyl-4-chlorobenzoate (500mg, 2.94mmol) and *N,N*-dimethylmethanesulphonamide (302mg, 2.45mmol) in ethylene glycol dimethyl ether (15ml) was added NaH (60% suspension in mineral oil, 265mg, 6.62mmol). The reaction was warmed to 85°C and stirred at this temperature for 3 hours. The reaction was cooled to room temperature and then quenched with water (~40ml). The water was extracted with ether then the ether was extracted with 1M NaOH. The aqueous fractions were combined and acidified to ~pH3 by the addition of concentrated HCl. The resulting suspension was extracted with DCM (2 x 40ml), the DCM layers were combined, washed with water and brine then dried, filtered and evaporated to yield an oil. This oil was purified by column chromatography (20g Si, DCM to 1% MeOH/DCM) to yield a solid (325mg, 42%). NMR (DMSO-*d*<sub>6</sub>): 2.95 (s, 6H), 4.55 (s, 2H), 7.50 (d, 2H), 8.00 (d, 2H); m/z 262 [M+H]<sup>+</sup>.


**Example 80**[2-(*N,N*-Diisopropylsulphamoyl)-2-(methyl)ethyl](4-fluorophenyl)ketone

Methyl iodide (51μl, 0.825mmol) was added to a stirred mixture of (*N,N*-diisopropylsulphamoylmethyl)(4-fluorophenyl)ketone (Example 3; 100mg, 0.33mmol) and potassium carbonate (114mg, 0.825mmol) in DMF (5ml) at room temperature under an inert atmosphere. The reaction mixture was stirred overnight before quenching with water (50ml) and then extraction with EtOAc (2x50ml). The organics were washed with brine (50ml) and then dried over magnesium sulphate. The solvent was then removed under reduced pressure and the resulting brown gum was purified by chromatography (eluting with 10% EtOAc/isohexane) to yield a colourless oil which solidified on scratching (57mg, 53%). NMR (DMSO-*d*<sub>6</sub>): 1.22 (d, 12H), 1.62 (s, 6H), 3.61 (m, 2H), 7.28 (m, 2H), 8.02 (m, 2H).



Examples 81-89

The procedure described in Example 80 was repeated using the appropriate starting material(s) in place of (*N,N*-diisopropylsulphamoylmethyl)(4-fluorophenyl)ketone (Example 3) to give the following Examples.




Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
81	2,4-Difluoro phenyl	i-Pr	i-Pr	1.36 (d, 12H), 1.65 (s, 6H), 3.72 (m, 2H), 6.87 (m, 2H), 7.90 (m, 1H)		Example 31
82	3-Chloro phenyl	Me	Me		290	Example 4
83	4-Chloro-3-fluoro phenyl	Me	Me		308	Example 20
84	4-Trifluoro-methyl phenyl	Me	Me	1.64 (s, 6H), 2.91 (s, 6H), 7.83 (m, 2H), 7.93 (m, 2H)	324	Example 17
85	4-Cyano phenyl	i-Pr	i-Pr	1.33 (d, 12H), 1.71 (s, 6H), 3.64 (m, 2H), 7.72 (m, 2H), 8.12 (m, 2H)		Example 38
86	4-Fluoro phenyl	Me	Pr		302	Example 9
87	4-Fluoro phenyl	Me	-(CH <sub>2</sub> ) <sub>2</sub> -OMe		318	Example 8
88	2,4-Difluoro phenyl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> -OMe	1.34 (d, 6H), 1.65 (s, 6H), 3.35 (s, 3H), 3.40 (m, 2H), 3.53 (m, 2H), 3.97 (m, 1H), 6.87 (m, 2H), 7.77 (m, 1H)	364	Example 61

Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	NMR	M/z	SM
89	4-Fluoro phenyl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> - O-iPr		374	Example 55

**Examples 90-92**

The procedure described in Example 80 was repeated using the appropriate starting material(s) in place of (*N,N*-diisopropylsulphamoylmethyl)(4-fluorophenyl)ketone (Example 53) to give the following Examples.



Ex	Ring A	R <sup>1</sup>	R <sup>2</sup>	M/z	SM
90	4-Fluorophenyl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> -O-iPr	358 (M-H) <sup>+</sup>	Example 55
91	2-Chlorothien-5-yl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> -OMe	352 (M-H) <sup>+</sup>	Example 60
92	Benzothien-2-yl	i-Pr	-(CH <sub>2</sub> ) <sub>2</sub> -O-iPr	396 (M-H) <sup>+</sup>	Example 54

**Example 93**10 [2-(*N,N*-Diisopropylsulphamoyl)ethyl][4-(*t*-butyldimethylsilyloxy)phenyl]ketone


Methyl iodide (2.33ml, 37.5mmol) was added to a stirred mixture of (*N,N*-diisopropylsulphamoylmethyl)[4-(*t*-butyldimethylsilyloxy)phenyl]ketone (Example 62; 7.5mmol) and potassium carbonate (5.18g, 37.5mmol) in acetone (60ml) at room temperature under an inert atmosphere. The reaction mixture was stirred overnight at room temperature before adding more methyl iodide (2.33ml, 37.5mmol) and heating at reflux for 1 hour. The reaction mixture was quenched with water (200ml) and then extracted with EtOAc (2x150ml). The organics were dried over magnesium sulphate before being removed under reduced pressure. The resulting orange oil was purified by chromatography (eluting with 10% EtOAc/isohexane) to yield a colourless oil (1.542g, 48%). NMR: 0.15 (s, 6H), 0.89 (s, 12H), 1.18 (m, 12H), 1.55 (d, 3H), 3.58 (m, 2H), 4.84 (q, 1H), 6.79 (d, 2H), 7.87 (d, 2H).

**Example 94**(*N,N*-Dimethylsulphamoylmethyl)(3-methylphenyl)ketone

*N,N*-dimethylaminomethanesulphonamide (37mg, 0.3mmol) and anhydrous THF (3ml) were placed in a tube. To this solution was added a 1M solution of lithium bis(trimethylsilyl)amide in THF (0.6ml, 0.6mmol). The reaction was allowed to stir at room temperature for 30 minutes. At this point a solution of ethyl 3-methylbenzoate (60mg, 0.36mmol) in anhydrous THF (1ml) was added. The reaction was stirred at room temperature for 2 hours then quenched with sat ammonium chloride solution (2ml). The tube was capped then shaken and allowed to settle. The organic layer was collected and evaporated under reduced pressure, the resulting crude material was purified by prep LCMS (1-40% over 9.5mins, acetonitrile/water, with a constant 5ml/min 4% formic acid / acetonitrile) to yield a solid (29mg, 40%). *M/z*: 242.

**Examples 95-105**

The procedure described in Example 94 was repeated using the appropriate ester in place of ethyl 3-methylbenzoate.



Ex	R <sup>1</sup>	NMR	MS
95	3-CH <sub>2</sub> =CHCH <sub>2</sub> -		268
96	4-CN		253
97	3-F	3.00 (s, 6H), 4.50 (s, 2H), 7.35 (m, 1H), 7.50 (m, 1H), 7.75 (m, 1H), 7.85 (d, 1H)	246
98	3-CN		253
99	4-MeC(O)-		268 (M-H) <sup>+</sup>
100	2-F		246
101	2,4-diF	2.95 (s, 6H), 4.60 (s, 2H), 6.90 (m, 1H), 7.00 (m, 1H), 7.95 (m, 1H)	262 (M-H) <sup>+</sup>
102	3-MeO, 4-Cl		290 (M-H) <sup>+</sup>

Ex	R <sup>1</sup>	NMR	MS
103	4-HC≡C-		252
104	4-CH <sub>2</sub> =CH-CH <sub>2</sub> -O-		284
105	3,4,5-triMeO-	NMR (DMSO-d <sub>6</sub> ): 2.80 (s, 6H), 3.80 (s, 3H), 3.85 (s, 6H), 4.95 (s, 2H), 7.35 (s, 2H)	318

### Preparation of Starting Materials

The starting materials for the Examples above are either commercially available or are readily prepared by standard methods from known materials. For example the following 5 reactions are illustrations but not limitations of the preparation of some of the starting materials used in the above reactions.


#### Method 1

##### 1-Methoxy-2-[N-(methyl)mesylamino]ethyl

10 To a stirred solution of N-(2-methoxyethyl)methylamine (750mg, 8.43mmol) and triethylamine (938mg, 9.27mmol) in anhydrous DCM (60ml) at 0°C was added mesylchloride (966mg, 8.43mmol). The reaction was stirred at 0°C for 10 minutes then allowed to warm to room temperature and left to stir for a further 30 minutes. The reaction was then transferred to a separating funnel and washed with 2M HCl (20ml), water (20ml) and brine (20ml) then  
15 dried, filtered and evaporated to yield the product as a pale yellow oil (935mg, 67%). NMR: 2.85 (s, 3H), 2.95 (s, 3H), 3.40 (m, 5H), 3.55 (t, 2H).

#### Methods 2-3

The procedure described in Method 1 was repeated using the appropriate amine in  
20 place of N-(2-methoxyethyl)methylamine.



Meth	R <sup>1</sup>	R <sup>2</sup>	NMR
2	Me	Pr	0.95 (t, 3H), 1.60 (m, 2H), 2.80 (s, 3H), 2.95 (s, 3H), 3.10 (t, 2H)
3	Et	i-Pr	1.25 (m, 9H), 2.85 (s, 3H), 3.20 (q, 2H), 4.10 (m, 1H)

4	H	$-(CH_2)_2-OCH_3$	2.99 (s, 3H), 3.31(m, 2H), 3.40 (s, 3H), 3.53 (m, 2H), 4.92 (br s, 1H)
5	H	$-(CH_2)_2-O-iPr$	1.17 (d, 6H), 2.99 (s, 3H), 3.28 (m, 2H), 3.55 (t, 2H), 3.62 (m, 1H), 4.82 (br s, 1H)

**Method 6**1-(*N,N*-Dimethylamino)-2-[*N*-(methyl)methylamino]ethyl


To a stirred solution of *N,N,N*-trimethylethylenediamine (1.02g, 10mmol) and triethylamine (1.11g, 11mmol) in anhydrous DCM (70ml) at 0°C was added mesylchloride (1.15g, 10mmol). The reaction was stirred at 0°C for 10 minutes then allowed to warm to room temperature and left to stir for a further 30 minutes. Volatiles removed under reduced pressure and resulting oil taken up in DCM (60ml) then washed with 2M NaOH (30ml) and brine (30ml). The solvent was removed under reduced pressure to yield the product as an oil (962mg, 53%). NMR: 2.30 (s, 6H), 2.50 (t, 2H), 2.85 (s, 3H), 2.90 (s, 3H), 3.30 (t, 2H).

**Method 7**Propane-2-sulphonic acid dimethyl amide

To a stirred solution of 2M dimethylamine in THF (5.35ml, 10.7mmol) and pyridine (865μl, 10.7mmol) in anhydrous THF (10ml) was added isopropylsulphonylchloride (1ml, 8.9mmol). The reaction was stirred at room temperature overnight. Volatiles removed under reduced pressure and resulting oil taken up in EtOAc (60ml) then washed with 0.5M HCl (30ml) and water (30ml). The solvent was dried over magnesium sulphate and then removed under reduced pressure to yield the product as an orange oil (688mg, 51%). NMR: 1.36 (d, 6H), 2.93 (s, 6H), 3.25 (m, 1H).

**Methods 8-9**

The procedure described in Method 7 was repeated using the appropriate amine in place of dimethylamine.



Meth	R <sup>1</sup>	R <sup>2</sup>	NMR
8	Me	i-Pr	(DMSO-d <sub>6</sub> ) 1.10 (d, 6H), 1.18 (d, 6H), 2.68 (s, 3H), 3.24 (m, 1H), 3.94 (m, 1H)
9	H	i-Pr	(DMSO-d <sub>6</sub> ) 1.05 (d, 6H), 1.10 (d, 6H), 3.06 (m, 1H), 3.40 (m, 1H), 6.80 (d, 1H)

**Method 10***N*-Isopropyl-*N*-(2-methoxy-ethyl)-methanesulphonamide


- 5 To a stirred solution of *N*-(2-methoxy-ethyl)-methanesulphonamide (Method 4) (1.2g, 7.7mmol) in DMF (20ml) under an inert atmosphere was added sodium hydride (400mg, 10mmol). 2-Bromopropane (1.73ml, 18.48mmol) was added and the reaction heated at 70°C for 7h. The reaction mixture was quenched with water (100ml) and then extracted into EtOAc (100ml). The organics were further washed with brine (100ml) before being dried over
- 10 magnesium sulphate. The solvent was then removed under reduced pressure to yield the product as a yellow oil (600mg, 40%). NMR: 1.26 (d, 6H), 2.88 (s, 3H), 3.31 (m, 2H), 3.38 (s, 3H), 3.55 (m, 2H), 4.09 (m, 1H).

**Method 11**15 *N*-Isopropyl-*N*-(2-isopropoxy-ethyl)-methanesulphonamide

The procedure described in Method 10 was repeated using *N*-(2-isopropoxyethyl)methanesulphonamide (Method 5) in place of *N*-(2-methoxy-ethyl)-methanesulphonamide. NMR: 1.16 (d, 6H), 1.27 (d, 6H), 2.90 (s, 3H), 3.29 (m, 2H), 3.56 (m, 3H), 4.09 (m, 1H).

Claims

1. The use of a compound of formula (I):



(I)

wherein:

Ring A is selected from carbocyclyl or heterocyclyl;

R<sup>1</sup> is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, N-(C<sub>1-4</sub>alkyl)amino, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, N-(C<sub>1-4</sub>alkyl)carbamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, N-(C<sub>1-4</sub>alkyl)sulphamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino, tri-(C<sub>1-4</sub>alkyl)silyloxy, carbocyclyl, heterocyclyl, carbocyclylC<sub>0-4</sub>alkylene-Y- and heterocyclylC<sub>0-4</sub>alkylene-Y-; wherein R<sup>1</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>6</sup>; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>7</sup>;

n is 0-3; wherein the values of R<sup>1</sup> may be the same or different;

R<sup>2</sup> and R<sup>3</sup> are independently selected from hydrogen, hydroxy, amino, cyano, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, N-(C<sub>1-4</sub>alkyl)amino, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, carbocyclyl, heterocyclyl, carbocyclylC<sub>1-4</sub>alkyl, heterocyclylC<sub>1-4</sub>alkyl; wherein R<sup>2</sup> and R<sup>3</sup> may be independently optionally substituted on carbon by one or more groups selected from R<sup>8</sup>; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>9</sup>;

one of R<sup>4</sup> and R<sup>5</sup> is selected from C<sub>1-4</sub>alkyl and the other is selected from hydrogen or C<sub>1-4</sub>alkyl; wherein R<sup>4</sup> and R<sup>5</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>10</sup>;

Y is -S(O)<sub>a</sub>-, -O-, -NR<sup>12</sup>-, -C(O)-, -C(O)NR<sup>13</sup>-, -NR<sup>14</sup>C(O)- or -SO<sub>2</sub>NR<sup>15</sup>-; wherein a is 0 to 2;

R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup> and R<sup>15</sup> are independently selected from hydrogen, phenyl and C<sub>1-4</sub>alkyl;

R<sup>6</sup> and R<sup>8</sup> are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl;

$C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ carbamoyl,  $C_{1-4}$ alkylS(O) $_a$  wherein  $a$  is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl  
 5 and heterocyclyl; wherein  $R^6$  and  $R^8$  may be independently optionally substituted on carbon by one or more  $R^{11}$ ;

$R^{10}$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ amino,  
 10  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ carbamoyl,  $C_{1-4}$ alkylS(O) $_a$  wherein  $a$  is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ sulphamoyl,  $C_{1-4}$ alkylsulphonylamino; wherein  $R^{10}$  may be independently optionally substituted on carbon by one or more  $R^{16}$ ;

$R^7$  and  $R^9$  are independently selected from  $C_{1-4}$ alkyl,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkylsulphonyl,  
 15  $C_{1-4}$ alkoxycarbonyl, carbamoyl,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl;

$R^{11}$  and  $R^{16}$  are independently selected from halo, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxyl, methylamino, ethylamino, dimethylamino,  
 20 diethylamino,  $N$ -methyl- $N$ -ethylamino, acetylamino,  $N$ -methylcarbamoyl,  $N$ -ethylcarbamoyl,  $N,N$ -dimethylcarbamoyl,  $N,N$ -diethylcarbamoyl,  $N$ -methyl- $N$ -ethylcarbamoyl, methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl, ethylsulphonyl, methoxycarbonyl, ethoxycarbonyl,  $N$ -methylsulphamoyl,  $N$ -ethylsulphamoyl,  $N,N$ -dimethylsulphamoyl,  $N,N$ -diethylsulphamoyl or  $N$ -methyl- $N$ -ethylsulphamoyl;


25 or a pharmaceutically acceptable salt thereof;

in the manufacture of a medicament for use in the inhibition of 11 $\beta$ HSD1.



ABSTRACTTITLE: CHEMICAL COMPOUNDS

5 Compounds of formula (I):



(I)

wherein variable groups are as defined within; for use in the inhibition of 11 $\beta$ HSD1 are described.

THE PATENT OFFICE

18 NOV 2003

Received in Patents  
International Unit